



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2014

**Erratum: Brightest cluster galaxies in cosmological simulations with
adaptive mesh refinement: successes and failures**

Martizzi, Davide ; Teyssier, Romain ; Moore, Ben

DOI: <https://doi.org/10.1093/mnras/stu1882>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-154155>

Journal Article

Published Version

Originally published at:

Martizzi, Davide; Teyssier, Romain; Moore, Ben (2014). Erratum: Brightest cluster galaxies in cosmological simulations with adaptive mesh refinement: successes and failures. Monthly Notices of the Royal Astronomical Society, 445(1):1002.

DOI: <https://doi.org/10.1093/mnras/stu1882>

Erratum: Brightest cluster galaxies in cosmological simulations with adaptive mesh refinement: successes and failures

by Davide Martizzi,¹★ Jimmy,² Romain Teyssier³ and Ben Moore³

¹Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

²George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA

³Institute for Computational Science, University of Zurich, CH-8057 Zurich, Switzerland

CORRECTION TO FIG. 7

The paper ‘Brightest Cluster Galaxies in Cosmological Simulations with Adaptive Mesh Refinement: Successes and Failures’ was published in MNRAS 443, 1500 (2014). We report here that a duplicate of Figure A1 was incorrectly used in place of Fig. 7 on page 1506.

The correct figure is provided below. This error has no consequences on the rest of the paper and its conclusions.

ACKNOWLEDGEMENTS

The simulations have been run on the Monte Rosa cluster at CSCS, Switzerland. DM is working as a postdoctoral fellow at University of California at Berkeley and his work is funded by the Swiss National Science Foundation.




Figure 7. Ellipticity (edge-on view) versus angular momentum probe parameter λ_R measured within the half-light radius. This plot is used to separate fast rotators and slow rotators; here, the separation is represented by the black solid line $\lambda_R = 0.33 \times \sqrt{\epsilon}$. Values measured within the effective radius of a sample of real BCGs from Jimmy et al. (2013) are also compared to our results (blue circles). The Jimmy et al. ellipticities are apparent and have not been corrected.

★ E-mail: dav.martizzi@berkeley.edu

This paper has been typeset from a \LaTeX file prepared by the author.